

# Impact of ecological conditions on population dynamic of *Phenacoccus solenopsis* Tinsley and associated predators in sweet potato fields ELBeheira Governorate, Egypt

Elsayed A. Refaei<sup>1\*</sup>, El-sarand E. A.<sup>1</sup>, Neveen E. El-Metwaly<sup>1</sup>, Abd El-Salam A. Farag<sup>1</sup>, Ekram A. Abdou<sup>1</sup> and Ismael MM<sup>1</sup>

<sup>1</sup> Plant Protection Research Institute, Agricultural Research Centre, Dokki, Egypt

Correspondence Author: Elsayed A. Refaei

Received 20 Feb 2024; Accepted 25 March 2024; Published 5 April 2024

## Abstract

The mealy bug, *Phenacoccus solenopsis* and the two spotted spider mite, *Tetranychus urticae* are a highly polyphagous sap sucking pests, without a hibernation period, as both are active all the year round. They are destructive pests of important vegetables and field crops. The present study was carried out at Rasheed region, Beheira Governorate during 2021 and 2022 seasons, to monitor the population fluctuations of mealybug and its predators; *Coccinella undecimpunctata* L., *Chrysoperla carnea* Steph, *Scymnus* spp. and *Hippodamia tredecimpunctata* in the first season, the infestation of *P. solenopsis* started on sweet potato plants at low numbers, the population increased gradually to reach its highest peak during 30<sup>th</sup> July (115.00 individual). This peak was followed by a relatively high population of the three predators, *C. undecimpunctata* (4.00 individual), *Ch. carnea* (8.00 individ.), *Scymnus* sp. (3.00 individ.) and *H. tredecimpunctata* (2.25 individ.). The second peak of the pest was detected on 21<sup>st</sup> Aug. (259.05 individual), synchronized with the peaks of the four, predators. During 2022, the first incidence of the mealybug was recorded on 30<sup>th</sup> July (114.97individ.). This peak was associated with the peaks of the four predators. The second peak of *P. solenopsis* was observed on 28<sup>th</sup> Aug. (196.50 individ.), it coincided with the peaks of the four predators. The infestation of *P. solenopsis* and *T. urticae* and their predators were highly significantly correlated with weather factors (Min.°c. and RH).

**Keywords:** *Phenacoccus solenopsis*, sweet potato, predators

## Introduction

The mealybug, *Phenacoccus solenopsis* Tinsley (Hemiptera: Pseudococcidae) is one of the most important insects causing severe economic losses to the yields, feeding on a wide variety of more plant species including vegetables, field crops, bushes, maize, weeds and ornamentals (Nagrare 2012; Fallahzadeh *et al.*, 2014; Abd razzik *et al.*, 2015; Refaei *et al.* 2016 and El-sarand 2018) [23, 12, 3, 25]. El-Zahi and Abd- Elsalam (2017) [10] and Hameed *et al.*, (2014) [16] found that the population density of the mealy bug was positively correlated with weather factors. Sweet potato *Ipomoea batatas* L crop is one of the food crops in the Egyptian economy for man and animal. Recently, sweet potato crop is infested with the newly introduced insect pests in Egypt, *P. solenopsis*. Economic damage was recorded on vegetable, cotton, maize and soybean, reached field death in sever conditions (Arif *et al.*, 2009, Bader *et al.*, 2020 and El-Zahi 2017 [4, 5, 8] and Abbas *et al.*, 2010) [1]. *P. solenopsis* was appeared for the first time in Egypt in September. Nabil and Hegab (2019) [22] indicated into positive correlation between *P. solenopsis* and temperature and significant negative correlation with relative humidity. El-Fakharany (2020) [9] showed that high population density of *P. solenopsis* was recorded in August and September and the infestation was high and positively correlated with the temperature and negative with relative humidity. In general, during the season, infestations of sweet potato fields, during the growth crop stage, cause severe economic losses to the yield. Ibrahim (2018) [18] found that five larvae of *Ch. carnea*/ 100 individual of the mealy bug can be

applied as a biological control with *P. solenopsis*. Farhan *et al.*, (2011) [13] found that the lacewing, *Ch. carnea* was more efficient as biological control against. Bakry *et al.*, (2023) [16] showed that *P. solenopsis* infestation in okra plants can be found on all plants parts five weeks after planting. Bader *et al.*, (2020) [5] recorded the pest for the first time in September in Egypt. noticed on sweet potato for many years it is an economic pest of various field crops in Egypt (Jakubwska and Fiedler, 2014) [20]. The information generated may be used for designing a comprehensive pest management program and prediction models for the cotton mealybug. The present study was planned to infestation the population fluctuation of *P. solenopsis* on sweet potato as affected by predators and to determine its, activity to certain synthetic predators under field conditions.

## Materials and methods

### a) Land preparation and sowing

This experiment was conducted carried out at Edfina Rasheed region, Beheira Governorate during 2021and 2022 summer seasons, in an area 2000 m<sup>2</sup>, divided into four equal parts. The land was prepared by laughing three times with calcium superphosphate at the rate of 250 kg /fed. Sweet potato (*Ipomoea batatas* L.) seedlings were Trans in the presence of water at the upper third of the furrows at the beginning of May in both years of study. Sulphur potassium (100 kg/fed.) and nitrogen fertilizer were app as recommended.

### b) Sampling for counting the predators

One month after sweet potato Trans planting, 25 plants were pulled out from each plot. The plants were gently confined in plastic bags and transferred to the laboratory of counting the four considered predators, *C. undecimpunctata*, *Ch. carnea*, *Scymnus sp.* and *Hippodamia tredecimpunctata*. This sampling technique was followed for four examinations, as the plants were still young with small size. After that, the sample was one branch of potato plants that was cut gently and confined in

plastic bags, as 25 branches were taken from each replicate to visually count the numbers of the three abovementioned predators. The sampling began on June 1<sup>st</sup> up to June 5<sup>th</sup>.

### c) Sampling for counting mealybug

Mealybug, *Phenacoccus solenopsis* Tinsly (Hemiptera: Pseudococcidae) was counted on 25 leaflets per replicated. The leaflets were weekly picked up and transferred to the laboratory for counting both arthropods, using binocular microscope.



**Fig (1):** Photographs showing the infestation by *P. solenopsis* on sweet potato plants

The current work depends on examining and identifying all the samples that were examined.

It was collected directly from the field and then transported to the laboratory where it was harvested. Learn about them in the Insect Survey and Classification section.

Plant Protection Research Institute of the Ministry of Agriculture Group [MAC];



**Fig (2):** Photographs showing of the predator, *Hippodamia tredecimpunctata* on sweet potato plants

## Results and discussion

### a) Population fluctuations of *Phenacoccus solenopsis*

In 2021 season, nymphs and adults of *P. solenopsis* were completely absent throughout June (Table 1). By the first week of July, the density of this mealybug was 29.75 nymphs and adults / 25 leaflets. The insect density gradually increased to exhibit the first peak (115.00 individuals / 25 leaflets) on July 30<sup>th</sup>. However, the highest peak (259.50 nymphs and adults/ 25 leaflets) was found on August 21<sup>st</sup>.

In 2022 season (Table 2), almost the insect population fluctuation was similar to those of the first season, with a peak of 114.97 nymphs and adults / 25 leaflets on Jul. 30<sup>th</sup>, with the highest peak (148.25 individuals) on August 28<sup>th</sup>. Overall mean

of the first season was higher than that of the second season; 79.29 and 71.02 nymphs and adults, respectively.

These results agree with Refai (2016) [25]. The mealy bug infestation appeared late in low numbers by late June up to mid-July during season 2016.

### b) Population fluctuations of predators

#### i) *Coccinella undecimpunctata*

In 2021 season, population densities of *C. undecimpunctata* were very low up to mid- June, and relatively increased by late June. Then, the predator population fluctuation to exhibited a small peak of 4.00 and adults /25 potato branches on Jul. 30<sup>th</sup>. In 2022, almost the same trend was detected, but with relatively high two peaks on July 30<sup>th</sup> and August 28<sup>th</sup>, with 6.75 and 5.75 predatory individuals, respectively. In a comparison, *C. undecimpunctata* individuals were relatively higher (5.75) in the second season than in the first one (3.50 adults/25 plants or branches).

#### ii) *Chrysoperla carnea*

In 2021, only two peaks of *C. carnea* with 8.00 and 6.75 larvae/25 potato branches were detected on July 30<sup>th</sup> and August 21<sup>st</sup>, respectively. In 2022 season, two peaks were found with 7.75 and 7.00 larvae/25 branches on July 30<sup>th</sup> and August 28<sup>th</sup>, respectively. Overall means in both seasons were similar.

#### iii) *Scymnus interruptus*

Overall mean of 2022 season was obviously higher (4.94 adults/ 25 branches) than that of 2021 season (3.86 adults). In 2021 season, the predator was observed with considerable high numbers by late August. In 2022 season, the insect densities were relatively high by late June, late August and early September. Refai The complex of insect predators and true spiders was detected in two peaks in each season.

**iv) *Hippodamia tredecimpunctata***

In 2021 season, population densities of *H. tredecimpunctata*, it appears late in July in small numbers and the population increases in August 21<sup>th</sup> (4.25 adults/25 plants or branches).

In 2022, It appears late in July in small numbers and the population increases in Jul 30<sup>th</sup> (5.25 adults/25 plants or branches).

**Table 1:** Averg numbers of Population fluctuation of *P. solenopsis* and associated predators on sweet potato branches, at Edfina- Rashed region in season 2021

Date of sampling	Av. No. per 25 plant				
	<i>Phenacoccus solenopsis</i>	<i>Coccinella undecimpunctata</i>	<i>Chrysoperla carnea</i>	<i>Scymnus interruptus</i>	<i>Hippodamia tredecimpunctata</i>
Jun., 1	0.00	0.00	0.00	0.00	0.00
8	0.00	1.75	2.75	1.50	0.00
15	0.00	1.25	2.75	1.75	0.00
22	0.00	3.75	2.00	2.50	0.00
30	0.00	3.25	3.75	1.75	0.00
Av. No. per 25 branches					
Jul.,7	29.75	2.75	4.00	2.25	1.25
15	35.75	2.00	3.50	3.00	1.75
22	48.25	2.50	2.75	2.25	1.75
30	115.00	4.00	8.00	3.00	2.25
Aug.7	55.00	3.50	4.75	3.75	2.25
14	110.25	3.25	5.25	4.25	1.25
21	259.50	3.25	6.75	11.75	4.25
28	201.50	3.50	4.00	13.75	1.00
Sept., 5	255.25	3.25	4.75	2.50	0.75
Overall ± SE	79.29±3.25	2.58±0.01	3.93±0.21	3.89±0.01	1.180 ±0.33

**Table 2:** Averg numbers of population fluctuation of *P. solenopsis* and associated predators on sweet potato branches, at Edfina- Rashed region in season 2022

Date of sampling	Av. No. per 25 plant				
	<i>Phenacoccus solenopsis</i>	<i>Coccinella undecimpunctata</i>	<i>Chrysoperla carnea</i>	<i>Scymnus interruptus.</i>	<i>Hippodamia tredecimpunctata</i>
Jun., 1	0.00	0.00	0.00	0.00	0.00
8	0.00	0.00	0.00	0.00	0.00
15	0.00	2.25	2.00	0.00	0.00
22	0.00	3.70	3.00	10.05	1.75
30	0.00	3.50	4.25	10.75	2.25
Av. No. per 25 branches					
Jul.,7	19.25	2.50	2.25	10.05	3.75
15	50.05	2.75	2.75	2.25	3.00
22	110.25	3.25	7.25	3.00	4.75
30	114.97	6.75	7.75	4.25	5.25
Aug.7	111.00	3.75	4.75	3.05	2.25
14	148.25	3.25	5.50	2.75	1.75
21	81.5	4.75	6.50	2.05	1.25
28	196.50	5.75	7.00	10.75	3.05
Sept., 5	162.52	3.50	4.25	10.25	2.25
Overall ± SE	71.02±3.12	3.26±1.01	4.13±0.20	4.94	1.18±0.32

**3. Correlations between populations of pests and their associated predators with some weather factors**

Data in Table (3) show that maximum temperature correlated with negative values, in most cases, with populations of *P. solenopsis* and their associated predators. However, the minimum temperatures and relative humidity were usually correlated with positive and highly significant values with the considered insect pests and their associated predators in the two

study seasons.

These results disagree with El-sarand *et al.*, (2018) [25]. They found that both temperature (max and min) and relative humidity had a negative and insignificant effect on the population *P. solenopsis* in the two seasons on soybean plants, while El- Fakhary (2020) [9] obtained that a positive significantly correlation between populations of mealy bug, in eggplant and okra, and each of minimum and maximum.

**Table 3:** Correlation coefficients between some weather factors and main pests and their predators on sweet potato plants

Factor	<i>Phenacoccus solenopsis</i>	<i>Coccinella undecim-punctata</i>	<i>Chrysoperla carnea</i>	<i>Scymnus sp.</i>	<i>Hippodamia tredecimpunctata</i>
2021					
Max. Tem(°c)	-0.246	+0.321	-0.184	-0.218	+0.352
Min. Tem (°c)	+0.359*	+0.567**	+0.454**	+0.462*	+0.403*
RH%	+0.445*	+0.689**	+0.500**	+0.354*	+0.533**
2022					
Max. Tem(°c)	+0.351	-0.211	-0.213	-0.189	-0.201
Min. Tem (°c)	+0.401*	+0.670**	+0.489**	+0.423*	+0.525**
RH%	+0.390*	+0.712**	+0.514**	+0.462*	+0.477*

\*significant,  $P \leq 0.05$  -\*\* highly significant,  $P \leq 0.01$

Temperature and relative humidity. Contracting to our results, Ramzan *et al* (2019) [24] observed that both *Chrysoperla sp* and coccinellids spp were active, in cotton fields in June and September, respectively, they also found significant correlations between both predators and *P. solenopsis*. Shahid *et al* (2022) [28] indicated that integrated pest management (IPM) is the best way to control *P. solenopsis*, and they added that biological control is an important compound for the control of scale insects. Hameed *et al* (2012) [15] concluded that *P. solenopsis* caused significant economic losses. The adults and immature stages of *T. urticae*, feeding on sweet potato, result in high levels of plant destruction (Muluken *et al* 2016) [21] damage of 103 million bales, resulting in very large economic damage.

## References

- Abbas G, Arif MJ, Ashfaq M, Aslam M and Shafqat S. Host plants distribution and overwintering on cotton mealybug, *Phenacoccus solenopsis*, (Hemiptera: Pseudococcidae). Int. J. Agric. Biol. 2010;12(3):421-425.
- AbdEL-Mageed AS Sanaa, Soad I, Abdel-Razak A, Haris HM. Ecological studies on the cotton mealybug, *Phenacoccus solenopsis* (Hemiptera: Pseudococcidae) on maize in Upper Egypt. Egypt. J. Plant Prot. Res. Inst. 2020;3(4):1098-1110.
- Abdel- Razzik KI, Attia AR, AbdelAziz M. Newly host plants of cotton mealybug, *Phenacoccus solenopsis* Tinsley (Hemiptera: Pseudococcidae) in Egypt. Egypt Acad. Biolog. Sci. 2015;8(3):31-33.
- Arif MI, Rafiq M, Ghaffar A. Host plants of cotton mealybug (*Phenacoccus solenopsis*): A new menace to cotton agro-ecosystem of Punjab, Pakistan. Int. J. of Agriculture and Biology. 2009;11:163-167.
- Bader AK, Al-Jboory IJ. First record of cotton mealybug, *Phenacoccus solenopsis* Tinsley 1898 (Hemiptera: Pseudococcidae) from Saudi Arabia. Bulletin Oepp/Eppo Bulletin. 2020;50(3):1-4.
- Bakry MMS, Badawy AMM, Lamiaa HYM. Spatial distribution and abundance of the mealybug, *Phenacoccus solenopsis* (Hemiptera: Pseudococcidae) infesting okra plants. Int. J. of Agric. Sci. 2023;5(3):1-17.
- Desai AV, Desai RDP, Bhandari GR. Population dynamic of mealybug, *Phenacoccus solenopsis* and its natural enemies on *Bt* cotton. The Pharmal Innovation J. 2022;11(7):1506-1512.
- El-Zahi ES, Farag AI. Population dynamic of *Phenacoccus solenopsis* Tinsley on cotton plants and its susceptibility to some insecticides in relation to the exposure method. Alexandria Science Exchange Journal. 2017;38(2):231-237.
- El-Fakharany SKM. Cotton mealybug *Phenacoccus solenopsis* (Hemiptera: Pseudococcidae) population density in eggplant and okra plantation and effect of some insecticides. Egypt. J. Plant Prot. Res. Inst. 2020;3(1):377-388.
- El-Zahi ES. Preferences and predatory potential of *Chrysoperla carnea* (Stephens) and *Coccinella undecimpunctata* Linnaeus on *Phenacoccus solenopsis* Tinsly: A New Threat to the Egyptian Economic Crops. Alexandria Sci. Exchange J. 2017;38(4):837-843.
- El-Zahi ES, Aref SA, Korish SKM. The cotton mealybug, *Phenacoccus solenopsis* Tinsly (Hemiptera: Pseudococcidae) as a new menace to cotton in Egypt and its chemical control. J. of Plant Prot. Res. Inst. 2016;56(2):111-115.
- Fallahzadeh M, Abdmaleki R, Saghaei N. Host plant of the newly invasive mealybug species, *Phenacoccus solenopsis* (Hemiptera: Pseudococcidae), in Homozgan province, Southern Iran. Entomo. Fauna. 2014;9(35):169-176.
- Farhan A, Akram W, Sajjad A, Iran AU. Management practice against cotton mealybug, *Phenacoccus solenopsis* Tinsley (Hemiptera:Pseudococcidae). Int. J. Agriculture and Biology. 2011;13(4):547-552.
- Gaber WM, Abou-elenien NF, Shehata EA. Population fluctuation of *Tetranychus urticae* on sweet potato (*Ipomoea batatas* L.) and effect of some insecticides. Egypt Acad. J. Biology. Sci. 2023;15(1):123-127.
- Hameed A, Aziz MA, Aheer GM. Impact of ecological conditions on biology of cotton mealybug, *Phenacoccus solenopsis* Tinsley (Steronorrhyncha: Coccoidea: Pseudococcidae) in laboratory. Pakistan J. of Zoology. 2012;44:685-690.
- Hameed A, Shahzad MS, Mehmood A, Ahmad S, Islam N. Forecasting and modeling of sucking insect complex of cotton under agro-ecosystem of Multan- Punjab, Pakistan. Pakistan J. of Agricultural Sciences. 2014;51(4):99-103.
- Hendawy AS, El-Fakharany SK, Hegazy FH, Samy MA. Survey and population fluctuation of arthropod pests and predators in sweet potato at Nile Delta, Egypt. Egyptian Acad. J. of Biolog. Sci. A. Entomol. 2017;10(7):277-285.
- Ibrahim SS. Study on cotton host plants of mealybug, *Phenacoccus solenopsis* (Tinsley) and efficiency of its controlling on cotton plants in Egypt release the predators *Chrysoperla carnea* (Stephens). J. Plant. and Path., Mansoura Univ. 2018;9(3):247-252.

19. Jaha RC, Bharpoda TM, Patel MG. Occurrence of mealybug, *Phenacoccus solenopsis* Tinsley *Phenacoccus solani* Ferris (Stemonorrhyncha: Coccoidea: Pseudococcidae) on cotton in Gujarat. Insect Environ. 2008;13(4):149-150.
20. Jakubwska M, Fiedler Z. Plantacje burak ow zagrozone przez prz, edziorki. Wa zny jest monitoring (Beet plantation endangered by mite) Poradnik Plantatora Buraka Cukrowego. 2014;2:53-54.
21. Muluken G, Dejene M, Kassaye A, Belay T. Red spider mite, *Tetranychus urticae* Koch (Arachnida: Acari: Tetranychadae): A threatening pest to potato (*Solanum Tuberosum* L.) production in eastern Ethiopia. Pest Management J of Ethiopia. 2016;19:53-59.
22. Nabil HA, Hegab MAM. Impact of some weather factors on the population density of *Phenacoccus solenopsis* Tinsley and its natural enemies. Egypt. Acad. J. Biolog. Sci. 2019;12(2):99-108.
23. Nagrare VS, Kumar R, Amutha M, Dharajothi B, Kranthi B, Vennila S, *et al.* A record of host plants of mealybug, *Phenacoccus solenopsis* Tinsley for devising ecofriendly management strategies. J. Ent. Res. 2012;36(4):327-344.
24. Ramzan M, Ullah UN, Ishiaq M, Murtaza G, Qayyum MA, Manzoor F. Population dynamic of natural enemies and their correlation with weather parameters in cotton. J. of Innovative Sci. 2019;5(1):40- 45.
25. Refai EA, El-Sarand EA, Khalif AA. Population fluctuation of certain sucking insects and their natural enemies associated with seed watermelon, *citrullus lanatus* at rasheed region, beheira governorate, Egypt. J. Plant Protection, Res. 2016;4(4):61-69.
26. Saif El-Deen UM, Gouda AEAI, Badawy AS. Effect of foliar spray with some micronutrients and slow release nitrogen fertilizers rates on productivity and quality of sweet potato (*Ipomoea batatas* L.). J. Plant Prod., Mansoura Univ. 2015;6(8):1277-1291.
27. Shahid MR, Arif MJ, Gogi MD, Javed N. Host plant preference and mortality analysis of *Phenacoccus solenopsis* in association with biochemical traits of different plant species. Int. J. of Agriculture and Biology. 2016;19(2):211-218.
28. Shahid MR, Ilahi F, Zahra T, Rehman HU, Qureshi MM, Kanwal S, *et al.* Host plant- mealybug (*Phenacoccus solenopsis*) interaction: A review. J. of Pure and Applied Agric. 2022;7(3):66-77.